



Colombian Journal of Anesthesiology

Revista Colombiana de Anestesiología

www.revcolanest.com.co

OPEN

 Wolters Kluwer

Base deficit, lactate clearance, and shock index as predictors of morbidity and mortality in multiple-trauma patients

Déficit de base, depuración de lactato e índice de choque como predictores de morbimortalidad en pacientes politraumatizados

Carlos Andrés Cortés-Samacá^a, Héctor Julio Meléndez-Flórez^a, Saúl Álvarez Robles^a, Eduardo Andrés Meléndez-Gómez^b, Carla Andrea Puche-Cogollo^c, Henry Jair Mayorga-Anaya^a

^a Medical School, Universidad Industrial de Santander (UIS), Bucaramanga, Colombia

^b Intensive Care Unit, Clínica Carlos Ardila Lulle, Floridablanca, Colombia

^c Post-Anesthesia Care Unit, Hospital Universitario de Santander (HUS), Bucaramanga, Colombia.

Keywords: Lactic Acid, Morbidity, Mortality, Multiple Trauma, Hemorrhage

Palabras clave: Ácido Láctico, Morbilidad, Mortalidad, Traumatismo Múltiple, Hemorragia

Abstract

Introduction: Physiological responses to hemorrhage are not proportional to blood loss. This has led to the use of scores such as the shock index (SI) or biomarkers such as lactate clearance (LC) and base deficit (BD) in the assessment of multiple trauma patients.

Objective: To determine the risk between no LC, persistence of BD, and severe SI at 6 hours, and postoperative morbidity and mortality.

Methods: Prospective cohort study with multiple trauma patients taken to surgery. SI, BD, and lactate levels were calculated on admission and after 6 hours; LC was estimated at 6 hours; and follow-up was performed after 28 days to determine morbidity and mortality. Inadequate LC was defined

as <20% at 6 hours, severe SI as >1, and abnormal BD as <-6 mmol/L.

Results: Overall, 196 patients were assessed. Morbidity and mortality were 46.24% and 19.69%, respectively. Relative risks for morbidity such as surgical reintervention, acute renal injury (ARI), pneumonia, need for vasopressors, and transfusions were significant for LC<20%. Severe SI and BD on admission were not found to be significant predictors, but after 6 hours, their behavior was similar to that of LC. In the final model, the significant variables for mortality were LC<20% at 6 hours, ARI, age >45 years, suture dehiscence, transfusions.

Conclusion: In multiple trauma patients in shock, low LC, severe SI, and persistent BD at 6 hours were significant for morbidity and mortality and their use as follow-up markers should be encouraged in resuscitation therapies for patients in shock.

How to cite this article: Cortés-Samacá CA, Meléndez-Flórez HJ, Robles SA, Meléndez-Gómez EA, Puche-Cogollo CA, Mayorga-Anaya HJ. Base deficit, lactate clearance, and shock index as predictors of morbidity and mortality in multiple-trauma patients. Rev Colomb Anestesiología. 2018;46:208-215.

Read the Spanish version of this article at: <http://links.lww.com/RCA/A107>.

Copyright © 2018 Sociedad Colombiana de Anestesiología y Reanimación (S.C.A.R.E.). Published by Wolters Kluwer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Correspondence: Departamento de Cirugía, Escuela de Medicina, Universidad Industrial de Santander, Carrera 32 No. 29-31, Bucaramanga, Colombia. E-mail: melendez@uis.edu.co

Rev Colomb Anestesiología (2018) 46:3

<http://dx.doi.org/10.1097/CJ9.0000000000000064>

Resumen

Introducción: Las respuestas fisiológicas a la hemorragia no son proporcionales a la pérdida de sangre, lo cual ha llevado a usar escalas como el índice de choque (IC) o biomarcadores como la depuración de lactato (LC) y déficit de base (DB) en la evaluación del paciente politraumatizado.

Objetivo: Determinar el riesgo entre no LC, persistencia DB e IC severo a las 6 horas y morbimortalidad postquirúrgica.

Métodos: Se realizó un estudio tipo cohorte prospectiva de pacientes politraumatizados, intervenidos quirúrgicamente. Se calculó LC a las 6 horas; el IC, DBEx y lactato al ingreso y control a las 6 horas; se realizó seguimiento a 28 días para determinar morbilidad y mortalidad. LCa inadecuado se definió como $<20\%$ a las 6 horas, IC severo: >1 y DB anormal: <-6 mmol/L.

Resultados: Evaluamos 196 pacientes. La morbilidad fue del 46.24% y la mortalidad del 19,69%. Los RR para morbilidad como reintervención quirúrgica, injuria renal aguda (IRA), neumonía, requerimientos de vasopresor y transfusiones fueron significativos para LC $<20\%$. El ICs y el DB severo al ingreso no fueron predictores significativos, pero a las 6 horas presentaron un comportamiento similar a la LC. En el modelo final, las variables significativas para mortalidad fueron: LC $<20\%$ en 6 horas, IRA, edad >45 años, dehiscencia, transfusiones.

Conclusiones: En pacientes politraumatizados en estado de choque, la baja LC, el IC y DB severo persistente, a las seis horas, fueron significativos para morbimortalidad y deben promoverse como marcadores de seguimiento en las terapias de reanimación en choque.

Introduction

Close to one-half of deaths due to trauma are related to hemorrhage, hence the need to recognize it early on to mitigate the effects of hypovolemic shock which leads to tissue hypoxia, anaerobic metabolism, and metabolic acidosis.¹ Vital signs and clinical scores have been traditionally used for recognizing shock states in patients with severe multiple trauma; however, various publications have shown that physiological responses to bleeding such as hypotension and tachycardia may not be proportional to the state of shock. This has created the need for using other predictors such as lactate clearance (LC), defined as a drop in lactate levels compared with baseline values; base deficit (BD),² defined as the amount of base required to bring the pH in 1L of blood to 7.4, as a buffer mechanism to maintain pH within normal limits; and shock index (SI), defined as the quotient between heart rate over systolic blood pressure,³ allowing to increase sensitivity in the identification of hypoperfusion states.⁴ Our objective was to assess the relative risk of each variable for morbidity and mortality.

Methods

Prospective analytical cohort study with a follow-up between April 2016 and July 2017 of patients diagnosed with hemorrhagic shock,⁵ defined as systemic arterial hypotension (SBP <90 mm Hg or mean arterial pressure (MAP) <70 mm Hg), associated with signs of tissue hypoperfusion and hyperlactatemia. Inclusion criteria were patients taken to emergency surgery who had the study variables: blood gases plus serum lactate, vital signs, use of blood products and vasopressors, and SI, identified on admission and after 6 hours. Exclusion criteria were patients who died intraoperatively or within the first 6 hours, and with no blood gases, electrolytes, or serum lactate measurements at 6 hours. LC was calculated using the formula⁶ (serum lactate T0 – serum lactate T6)/serum lactate T0 $\times 100$, where T0 was the initial lactate measurement and T6 lactate measurement at 6 hours, and it was considered adequate when values were lower than 20% as compared with the baseline measurement. Excess BD (-ExBD) was considered severe if <-6 mEq/L⁷; SI was calculated and classified as severe if the value was >1 ⁸; lactate, SI, and BD values were recorded on admission and after 6 hours. Follow-up was conducted after 28 days to determine mortality and morbidity, including pneumonia, surgical site infection (SSI), acute renal injury (ARI), suture dehiscence (including evisceration), and reintervention. In addition, the use of blood products and vasopressors was also recorded.

Relative risk and 95% confidence interval (RR and 95% CI) were calculated for morbidity and mortality for each variable, and managed with dichotomically. Univariate and bivariate analyses were performed, together with a binomial logistic regression. Finally, modeling and goodness of fit were done to assess for correlation between data found and expected data. Data analysis was performed using the Stata 14 software package. (StataCorp. 2015. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP.) A P value <0.05 was considered significant. The study was approved by the interinstitutional Universidad Industrial de Santander (UIS) -Hospital Universitario de Santander (HUS) ethics committee.

Results

Of a total of 380 patients in the target population, 196 met the inclusion criteria. Mean age was 30 years, 8.16% were female, 89.8% were classified as American Society of Anesthesiologists (ASA) 4, and 10.2% as ASA 5. The most frequent injury (40%) was penetrating chest injury (Table 1).

Transfusion therapy was used in 45.4% of the patients, and vasopressor therapy was used in 32%. Although it was up to anesthetist to judge which therapy to use, it is important to highlight that the institutional protocol in this type of patient is to use the transfusion kit (5IU of red

Table 1. General patient characteristics and type of trauma

Variable	Total	Open		Closed
		Stab wound	Gunshot wound	
Male n (%)	180 (91.84)	116 (59.2)	39 (19.9)	25 (12.76)
ASA 4 n (%)	176 (89.75)	117 (59.7)	36 (18.36)	23 (11.73)
ASA 5 n (%)	20 (10.25)	10 (5.1)	5 (2.5)	5 (2.5%)
Age	30 (18–89)*	29.9 (18–56)	28.27 (18–82)	39.5 (18–89)
Trauma distribution	% (Fr) n=196 n(%)	Stab wound n=128 n(%)	Gunshot wound n=41 n(%)	% (Fr) n=27 n(%)
Brain	17 (8,6)	0	0	17 (8,6)
Neck	12 (6,1)	11 (5,6)	1 (0,5)	0
Chest	80 (40,8)	69 (35,2)	8 (4,08)	3 (1,5)
Abdomen	73 (37,2)	37 (18,8)	29 (14,7)	7 (3,5)
Extremities	14 (7,1)	11 (5,6)	3 (1,5)	0

* Age in average of years (minimum - maximum).
 ASA: American Society of Anesthesiologists physical status classification.
 Source: Authors.

blood cells, 5IU of platelets and 5IU of plasma). The interquartile range for length of stay was 7 days, and 12 days in the intensive care unit (ICU). Morbidity at 28 days was 46.24%, with a predominance of surgical reintervention and ARI. Mortality was 9.69% (Table 2).

Behavior of excess base, shock index, and lactate clearance on admission and at 6 hours

The behavior of -ExBD, SI, and aLC within the first 6 hours was very variable. The drop in these values was greater for severe SI (sSI), and only 6.63% did not achieve normal values during that time period. -ExBD dropped by 45%, but only 59% showed non-critical values. After 6 hours, 70% persisted with a lactate level of >2mmol/L, lactacidemia was found in 20%, and aLC >20% was found in 84% of patients (Table 3).

Relative risk of morbidity according to sSI, aLC, and -ExBD

General morbidity was 46.24%, with predominance of reintervention and renal injury in 32%. In terms of associated variables, -ExBD on admission was not associated with morbidity, but its persistence at 6 hours was associated with high and significant RR, except for suture dehiscence. sSI behavior on admission was a risk factor for needing transfusion therapy, and there was a significant association between the persistence of this score at 6

hours and morbidity, except for pneumonia, dehiscence, and SSI. Inadequate LC at 6 hours was associated with significantly high RR for all causes of morbidity studied. sSI and -ExBD at 6 hours were associated with the highest risk of ARI (Table 4).

Transfusion therapy and use of vasopressors

RR for the need of transfusion and use of vasopressors was significantly higher for sSI, LC, and -ExBD only at 6 hours (Table 5).

Relative risk for mortality: SI, LC, and -ExBD

At 6 hours, hyperlactatemia, LC <20%, and sSI were associated with an elevated and significant RR for mortality, but the same was not found for -ExBD. These findings will be subject for discussion (Table 6).

Binomial model for mortality

The group was evaluated as a whole and binomial regression and modeling were performed assessing those variables that could behave as risk factors. It was found that age >45 years, transfusion of blood products, renal failure, and clearance lower than 20% 6 hours after admission were significant predictors of mortality with quite a high RR for acute renal failure (RR 49.19). Goodness

Table 2. Intraoperative, in-hospital, morbidity, and mortality variables

Variable	
Therapies employed	n (%)
Need for transfusion	89 (47.8)
Use of vasopressors*	59 (31,7)
Length of stay	Average (minimum maximum)
ICU stay (days)	1.59 (2–28)
Ward stay (days)	7.66 (1–28)
Mortality	IC 95% (%)
General (at 28 days)	19 (9.7)
Early (24 hours)	10 (5.1)
Morbidity	n=86 (46,24) n (%)
Reintervention	33 (17.7)
Acute renal injury	27 (14.5)
Pneumonia	12 (6.4)
Surgical site infection	11 (5.9)
Suture dehiscence	3 (1.6)

*Vasopressors: norepinephrine equivalent dose >0.1 µg/kg/minute and/or dopamine 5 µg/kg/minute).

†Need for transfusion: trauma kit (5U of platelets, 5U of red blood cells, 5U of plasma).

Source: Authors.

of fitness (Hosmer–Lemeshow) showed perfect fit between observed and expected results ($P=0.86$; deviance=35.76266) (Table 7).

Discussion

One of the first authors to document the association between persistent and morbidity and mortality was Abramson, who in 1993 found that mortality was higher in patients with delayed drop in lactate levels down to baseline after 24 to 48 hours, and that survival was higher among those who achieved 100% LC within the first 24 hours.⁹ Our study found that mortality increases with persistence of lactacidemia over periods of time as short as 6 hours. In the 1990s, it was shown that not only high serum lactate levels upon admission but also clearance within the first 24 hours^{10,11} reduced morbidity and mortality, which was observed in our study as the highest rates of complications and mortality were found in patients with LC < 20%. Research has emerged showing that initial monitoring is quite useful for predicting outcomes and that the trend toward serial monitoring is a valuable approach for in-hospital mortality.¹² In our study, the measurement of hyperlactatemia on admission was only associated with the risk of transfusion, and no association was found with mortality or other morbidities, unlike the LC rate LC lactate level at 6 hours, in which statistically significant associations were found. Among several studies, it is worth highlighting the systematic review by Kruse¹³ which found 8 retrospective studies of moderate quality, as well as a retrospective study by Odom et al,¹⁴ world authorities in perfusion and microcirculation, that present initial lactate and LC at 6 hours as independent predictors of mortality in trauma patients; the authors recommend conducting prospective studies.

Table 3. Lactate, excess base, and shock index values on admission and at 6 hours

Variable	Values on admission		Values at 6 hours	
	n (%)	Mean (min–max) SD	Mean (min–max) SD	n (%)
Lactate between				
2–4 mmol/L	51 (26.02)	3.52 (2.8–4) 0.38	2.19 (0.7–4) 0.88	101 (51.5)
Lactacidemia: Lactate >4 mmol/L	145 (74)	7.43 (4.2–19.1) 3.37	7.34 (4.2–18.1) 3.64	38 (19.3)
Lactate normality <2 mmol/L		—	1.4 (0.7–2) 0.40	57 (29.08)
Lactate clearance <20%		—	46.8% (–59.5–91) 27.15	32 (16.3)
Severe excess base deficit (<–6)	25 (12.7)	–8.4 (–27.3–1.2) 4.6	–3.9 (23.6–10) 4.8	81 (41.3)
Severe shock index (>1)	183 (93.3)	1.48	0.8	32 (16.3)

Min = minimum; Max = maximum; SD = standard deviation.

Source: Authors.

Table 4. Relative risk of morbidity on admission and at 6 hours according to excess base deficit, lactate clearance, and shock index

	-ExBD1		-ExBD		SI1		SI6		LC at 6 hours	
	RR (95% CI)	P	RR (95% CI)	P	RR (95% CI)	P	RR (95% CI)	P	RR (95% CI)	P
Surgical reintervention	2.26 (0.57-8.9)	0.2	7.95 (3.2-19.7)	<0.001	2.09 (0.31-14.1)	0.41	4.97 (2.8-8.8)	<0.001	7.22 (4.1-12.8)	<0.001
Transfusion therapy	1.71 (0.89-3.3)	0.061	2.09 (1.5-2.9)	<0.001	5.70 (2.9-37.5)	0.008	2.30 (1.8-2.92)	<0.001	1.85 (1.4-2.5)	<0.001
Surgical site infection	1.46 (0.19-10.9)	0.7	6.38 (1.4-28.8)	0.005	3.73 (1.2-11.7)	0.024	1.98 (0.55-7.1)	0.28	23.95 (5.4-105.6)	<0.001
Acute renal injury	0.84 (0.31-2.2)	0.72	36.91 (5.1-66.5)	<0.001	0.54 (0.17-1.43)	0.21	7.69 (3.9-14.9)	<0.001	12.64 (6.1-26.3)	<0.001
Need for vasopressors	0.76 (0.41-1.4)	0.42	4.49 (2.5-8.1)	<0.001	0.73 (0.31-1.7)	0.49	4.67 (3.1-7.04)	<0.001	5.32 (3.6-7.9)	<0.001
Pneumonia	0.43 (0.12-1.5)	0.18	7.09 (1.6-31.5)	0.002	1.90 (1.05-3.47)	0.034	1.98 (0.55-7.1)	0.28	7.45 (2.5-21.97)	<0.001
Suture dehiscence	1*	0.208	2.83 (0.3-30.8)	0.36	1.81 (0.47-7.1)	0.391	2.64 (0.24-28.3)	0.40	2.7 (0.24-28)	0.418

CI=confidence interval; -ExBD1=excess base deficit on admission; ExBD 6=excess base deficit at 6 hours; LC=lactate clearance at 6 hours; RR=relative risk; SI1=severe shock index on admission; SI6=severe shock index at 6 hours.

*All patients with dehiscence showed severe EB.

Source: Authors.

In a systematic review, González et al¹⁵ conclude that the LC value in trauma has not been standardized and that there is heterogeneity in the different study populations and their management. We propose that a cutoff point of 20% for LC is adequate given that higher LC percentages might be related with more aggressive fluid therapies and potential adverse events.¹⁶ In a prospective observational

cohort study,⁶ Regnier considers that early LC, defined as 2 and 4 hours, is found to be an important independent and prognostic variable, but with very close measurement periods. Given the dynamics of our services, patient volumes, and our teaching environment, we consider these periods to be very short and non-conductive to a cost-effective assessment of these patients.

Table 5. Relative risk of the need for transfusion and vasopressors according to severe shock index, severe base excess deficit, and lactate clearance

Variable	Transfusion therapy*		Vasopressor therapy**	
	RR (95% CI)	P	RR (95% CI)	P
Severe shock index on admission	5.70 (2.86-37.48)	0.0082	0.73 (0.31-1.70)	0.49
Severe shock index at 6 hours	2.30 (1.81-2.92)	<0.001	4.67 (3.10-7.04)	<0.001
Severe excess base deficit on admission (<-6)	1.71 (0.89-3.27)	0.06	0.76 (0.41-1.43)	0.42
Severe excess base deficit at 6 hours (<-6)	2.09 (1.52-2.86)	<0.001	4.49 (2.50-8.05)	<0.001
Lactate clearance <r 20% at 6 hours	1.85 (1.40-2.45)	<0.001	5.32 (3.56-7.94)	<0.001

CI=confidence interval; RR=relative risk.

* > 2 units of blood products.

** norepinephrine dose >0.1 mcg/kg/min or dopamine mcg/kg/min.

Source: Authors.

Table 6. Relative risk of mortality on admission and at 6 hours

Variable	Mortality RR (95% CI)			
	Admission		At 6 hours	
	RR (95% CI)	P	RR (95% CI)	P
Hyperlactatemia	2.98 (0.71–12.49)	0.10	5.92 (2.28–15.39)	<0.001
Severe excess base deficit (<-6)	1.24 (0.30–5.05)	0.75	6.16 (2.5–15.4)	0.000
Lactate clearance <20% at 6 hours	-		14.9 (5.78–38.39)	<0.001
Severe shock index	0.32 (0.11–0.97)	0.05	19.83 (7.05–55.79)	<0.001

CI=confidence interval; RR=relative risk.
Source: Authors.

More recently, Oliveros et al published a prospective cohort of multiple trauma patients admitted to the ICU in whom they measured lactate at 12 and 24 hours. Their study showed lactate behavior similar to that found in other publications, with an association between mortality and persistently elevated lactate levels greater than 2.35 at 24 hours.¹⁷ The study has the drawback of not including an assessment of intraoperative behavior or intraoperative complications; it also has a selection bias as it only assesses the most critically ill patients transferred to the ICU and does not take into consideration patients who came in a state of shock but had a good therapeutic response and completed their course in the Post-Anesthesia Care Unit (PACU) and/or general wards. Our study has the advantage of assessing patients from the time of admission to the institution and following up on morbidity and mortality at 28 days, a period that is long enough to identify the various complications occurring in these patients.

In 2016, Dekker et al¹⁸ compared the different methods for LC calculation, and initial lactate measurement and concluded that there is no superiority of LC calculation over initial lactate measurement in terms of predicting

morbidity and mortality. The author of this retrospective study mentions a potential selection bias as part of the limitations. In our study, we documented that LC is better able to predict morbidity and mortality as compared with initial lactate measurement.

As far as BD is concerned, Ibrahim et al,¹⁹ in a recently published systematic review of studies of the past 25 years, found that BD values >6 mmol/L are associated with severe injuries and complications such as acute respiratory distress syndrome (ARDS), acute lung injury, ARI, multiple organ failure, and increased mortality. In our study, we found an association with morbidities such as pneumonia, ARI, surgical reintervention, need for transfusion therapy, and vasopressors, although no correlation was found with the risk of mortality. Moreover, this review found that improvement of BD values within the first 48 hours was associated with a lower mortality rate, but we were unable to document that reduction in relation to BD values on admission or at 6 hours.

Mutschler et al identified that BD -6 mmol/L on admission is an adequate cutoff point for mortality, in contrast with our study, which did not find that correlation. That same study established a new classification for hypovolemic shock based on the BD value, finding a better relationship between the degree of shock and the need for transfusion when compared with the classification proposed in the Advanced Trauma Life Support. This is supported by a retrospective review of 2954 patients¹² which documented that, in 72% of patients with BD \geq -6 mmol/L, red blood cell transfusion was needed within the first 24 hours; this is also consistent with our results that show that patients with severe persistent BD at 6 hours are at twice the risk of needing transfusion therapy.

Finally, we discuss the use of the sSI. In a systematic review, Olaussen⁸ concluded that the sSI is simple, reproducible, and useful for predicting the need for transfusion, and observed that the \geq 1.0 threshold had a higher specificity, a reason why we also defined the value of >1.0 as the cutoff

Table 7. Binomial model for mortality

Variable	Relative risk (95% CI)	P
Acute renal injury	49.19 (6.82–354.51)	<0.001
Age >45 years	2.45 (1.64–3.65)	<0.001
Suture dehiscence	1.72 (1.15–2.56)	0.007
Lactate clearance <20% at 6 hours	1.42 (1.42–1.43)	<0.001
Transfusion therapy	1.17 (1.17–1.73)	<0.001

CI=confidence interval.
Source: Authors.

point. Our results also showed that an SI higher than 1 is predictive of the need for transfusion therapy.

In our setting, Charry³ published in 2015 a cohort study with 170 patients and found that an SI > 0.9 is predictive of worse prognosis at 24 hours after trauma. Another finding highlighted in that study as well as in ours is that there is a statistically significant correlation between serum lactate and an SI greater than 1 for mortality. However, our study has the additional advantage of not only predicting mortality with the SI but also predicting morbidity and the need for transfusion therapy and vasopressors.

In a retrospective cohort of 1419 patients, Mitra et al²⁰ found that patients receiving a load of 1L of crystalloids who persisted with a high SI had a higher need for transfusion therapy, higher mortality, and worse outcomes than patients who responded initially to fluid therapy. Although we did not assess specific therapeutic effects, we documented that if the SI does not improve within 6 hours, there is a dramatic increase in mortality, the need for transfusion and vasopressors, and morbidity.

In another retrospective cohort of 16,269 patients, McNab et al studied the relationship between severe pre-hospital SI and the length of hospital stay, length of stay in the ICU, number of days on mechanical ventilation, and the use of blood products; they found that an SI > 0.9 indicates a higher probability of transfer to the ICU, emergency surgery, or death.²¹ In our study, we determined the initial SI which, extrapolating to this study, would be equivalent to the pre-hospital SI, and we were only able to find a greater need for blood products. In terms of mortality prediction, we found that a severe SI on admission is a protective factor, with statistical significance. We believe that when these patients arrive in these conditions, the trauma teams react faster and applies damage control principles, leading perhaps to a better impact on morbidity and mortality.

In the same cohort, McNab et al discuss their concern regarding the accuracy of the SI in the elderly. In a subgroup analysis, correlations for patients between 16 and 60 years of age were positive ($P < 0.05$). In patients over 80, none of the correlations with the outcome variables were statistically significant. In patients over 60 years of age, there was a correlation between a higher SI and an increase in mortality rates, and the authors concluded that the accuracy of pre-hospital SI declines in patients over 60.²² In our study, the mean age of the population was 30 years. Consequently, outcomes have to be interpreted with caution in geriatric ages as physiological compensatory mechanisms are diminished in this population group when exposed to trauma. Moreover, other interventions such as the use of beta blockers, cardiac devices such as pacemakers, implantable cardiac defibrillators, hip surgery, and heart transplant prevent compensation and will skew presentation toward a less compromised index.²³ For this reason, lower cutoff points are proposed for this age group.

In terms of transfusion therapy, various studies have shown the association between transfusion of blood products and greater survival.^{24,25} Our finding that a severe SI on admission behaves as a protective factor may be secondary to the fact that patients arriving to our institution in such critical conditions trigger a protocol of massive transfusion, damage control surgery, and other resuscitation measures that impact mortality. If this index remains abnormal despite these measures, all compensatory mechanisms are overridden, and it becomes a statistically significant predictor of mortality. Other studies show that there is no reduction in mortality with the early use of transfusion therapy.²⁶ In contrast, there are no studies in humans supporting the use of vasopressors in resuscitation of multiple trauma patients and, actually, their early use for hemodynamic support in patients with hemorrhagic shock may be deleterious and has been associated with higher mortality.²⁷⁻²⁹ A meta-analysis of randomized clinical trials and observational studies comparing the different fluid administration strategies in trauma³⁰ concludes that initial resuscitation strategies using liberal fluid administration may be associated with higher mortality. However, the authors of this review report that the available studies are subject to a high risk of selection bias and have a high clinical heterogeneity; consequently, this result must be interpreted with caution.

Because of its design, analysis, and follow-up, our study has validity and an adequate level of evidence. Therefore, the results are valid and applicable to our study population.

Limitations

Although the sample size in this prospective study, as compared with other international series, could be considered a limitation, there was an initial pilot study used to calculate the sample size. Consequently, the results obtained are statistically significant and not only totally applicable to our Colombian population and to our institution but also valid for extrapolation to different geographic settings. Although the mean age of the patients in our study was 30 years, it was not possible to conduct adequate subgroup assessments to determine behavior at different stages of life. Neither comorbidities that could affect LC (liver or kidney failure) nor renal or respiratory abnormalities that could affect acid-base balance before trauma were identified. Although we documented the risk of mortality for transfusion therapy and the use of vasopressors, we did not stratify groups based on the type of resuscitation. Therefore, these results must be interpreted with caution as close to 90% of our institutional staff members follow the protocol for trauma resuscitation and transfusion therapy concomitantly with the use of vasopressors such as noradrenaline and vasopressin.

Conclusion

LC < 20% at 6 hours is a risk factor for mortality and morbidity. No statistically significant association between morbidity and mortality and severe BD on admission, but severe BD at 6 hours is a risk factor for mortality and morbidity, except for suture dehiscence. Severe SI on admission acts like a protective factor for mortality, but persistence at 6 hours is associated with higher mortality and morbidity, except for pneumonia, suture dehiscence, and SSI. Given that the methods (biomarkers: lactate and base excess; scale CI) used in this study are independent predictors of mortality, they must be taken into consideration to guide resuscitation therapy and to use hemodynamic target-guided therapy guided in patients showing no improvement of control measures at 6 hours.

Ethical responsibility

Human and animal protection. The authors declare that no human or animal experiments were performed for this research.

Data confidentiality. The authors declare that they followed the protocols of their institutions regarding patient data disclosure.

Privacy and informed consent. No patient data appear in this article.

Funding

The authors have not received any form of financial support or grant during the development of this manuscript.

Conflict of interest

The authors declare having no conflict of interest.

References

- Raux M, Le Manach Y, Gauss T, et al. Comparison of the prognostic significance of initial blood lactate and base deficit in trauma patients. *Anesthesiology* 2017;126:522–533.
- Saad S, Mohamed N, Moghazy A, et al. Venous glucose, serum lactate and base deficit as biochemical predictors of mortality in patients with polytrauma. *Ulus Travma Acil Cerrahi Derg* 2016;22:29–33.
- Charry JD, Bermeo JM, Montoya KF, et al. Shock index as a predictor of mortality in patients with penetrating trauma to the thorax. *Rev Colomb Cir* 2015;30:24–28.
- Paladino L, Sinert R, Wallace D, et al. The utility of base deficit and arterial lactate in differentiating major from minor injury in trauma patients with normal vital signs. *Resuscitation* 2008;77:363–368.
- Vincent JL, De Backer D. Circulatory Shock. *N Engl J Med* 2013;369:1726.
- Régnier MA, Raux M, Le Manach Y, et al. Prognostic significance of blood lactate and lactate clearance in trauma patients. *Anesthesiology* 2012;117:1276–1288.
- Callaway DW, Shapiro NI, Donnino MW, et al. Serum lactate and base deficit as predictors of mortality in normotensive elderly blunt trauma patients. *J Trauma* 2009;66:1040–1044.
- Olaussen A, Blackburn T, Mitra B, et al. Review article: shock index for prediction of critical bleeding post-trauma: a systematic review. *Emerg Med Australas* 2014;26:223–228.
- Abramson D, Scalea TM, Hitchcock R, et al. Lactate clearance and survival following injury. *J Trauma* 1993;35:8–584.
- Dellinger RP, Levy MM, Rhodes A, et al. Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med* 2013;39:165–228.
- Okello M, Makobore P, Wangoda R, et al. Serum lactate as a predictor of early outcomes among trauma patients in Uganda. *Int J Emerg Med* 2014;7:20.
- Mutschler M, Nienaber U, Brockamp T, et al. Renaissance of base deficit for the initial assessment of trauma patients: a base deficit-based classification for hypovolemic shock developed on data from 16,305 patients derived from the TraumaRegister DGU®. *Crit Care* 2013;17:R42.
- Kruse O, Grunnet N, Barfod C. Blood lactate as a predictor for in-hospital mortality in patients admitted acutely to hospital: a systematic review. *Scand J Trauma Resusc Emerg Med* 2011;19:74.
- Odom SR, Howell MD, Silva GS, et al. Lactate clearance as a predictor of mortality in trauma patients. *J Trauma Acute Care Surg* 2013;74:999–1004.
- González M, Morales CH, Sanabria Á. Lactate clearance as a prognostic factor in patients with penetrating trauma. *Rev Colomb Cir* 2016;31:34–43.
- Baxter J, Cranfield KR, Clark G, et al. Do lactate levels in the emergency department predict outcome in adult trauma patients? A systematic review. *J Trauma Acute Care Surg* 2016;81:555–566.
- Oliveros-Rodríguez H, Estupiñán-López R, Rodríguez-Gómez J. Lactate serial measurements and predictive validity of early mortality in trauma patients admitted to the intensive care unit. *Rev Colomb Anestesiología* 2017;45:166–172.
- Dekker SE, de Vries HM, Lubbers WD, et al. Lactate clearance metrics are not superior to initial lactate in predicting mortality in trauma. *Eur J Trauma Emerg Surg* 2016;42:1–11.
- Ibrahim I, Chor WP, Chue KM, et al. Is arterial base deficit still a useful prognostic marker in trauma? A systematic review. *Am J Emerg Med* 2016;34:626–635.
- Mitra B, Fitzgerald M, Chan J. The utility of a shock index ≥ 1 as an indication for pre-hospital oxygen carrier administration in major trauma. *Injury* 2014;45:61–65.
- McNab A, Burns B, Bhullar I, et al. A prehospital shock index for trauma correlates with measures of hospital resource use and mortality. *Surgery* 2014;152:473–476.
- McNab A, Burns B, Bhullar I, et al. An analysis of shock index as a correlate for outcomes in trauma by age group. *Surgery* 2013;154:384–387.
- Neideen T, Lam M, Brasel KJ. Preinjury beta blockers are associated with increased mortality in geriatric trauma patients. *J Trauma* 2008;65:1016–1020.
- O'Reilly DJ, Morrison JJ, Jansen JO, et al. Prehospital blood transfusion in the en route management of severe combat trauma: a matched cohort study. *J Trauma Acute Care Surg* 2014;77 (3 suppl 2):S114–S120.
- Shackelford SA, Del Junco DJ, Powell-Dunford N, et al. Association of prehospital blood product transfusion during medical evacuation of combat casualties in Afghanistan with acute and 30-day survival. *JAMA* 2017;318:1581–1591.
- Huang GS, Dunham CM. Mortality outcomes in trauma patients undergoing prehospital red blood cell transfusion: a systematic literature review. *Int J Burns Trauma* 2017;7:17–26.
- Sperry JL, Minei JP, Frankel HL, et al. Early use of vasopressors after injury: caution before constriction. *J Trauma* 2008;64:9–14.
- Plurad DS, Talving P, Lam L, et al. Early vasopressor use in critical injury is associated with mortality independent from volume status. *J Trauma* 2011;71:565–570.
- Gupta B, Garg N, Ramachandran R. Vasopressors: do they have any role in hemorrhagic shock? *J Anaesthesiol Clin Pharmacol* 2017;33:3–8.
- Wang CH, Hsieh WH, Chou HC, et al. Liberal versus restricted fluid resuscitation strategies in trauma patients: a systematic review and meta-analysis of randomized controlled trials and observational studies. *Crit Care Med* 2014;42:954–961.