



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Questions and Answers[☆]

Preguntas y respuestas

Fernando Raffan Sanabria^{a,b,*}

^a Anesthesiologist and Intensivist, University Hospital Fundación Santa Fe de Bogotá, Colombia

^b Professor, Universidad el Bosque and Universidad de los Andes, Bogotá, Colombia

This section includes questions prepared on the basis of the contents of the articles published in volume 42, number 1 of the *Colombian Journal of Anesthesiology*. Accept the challenge to test your reading comprehension and knowledge.

1. The reported incidence of postoperative laryngopharyngeal symptoms following general anesthesia is between 5 and 70% of the patients. Which of the following distractors is not associated with an increased occurrence thereof?
 - a. Use of succinylcholine
 - b. Male gender
 - c. Higher incidence with endotracheal tube versus laryngeal mask
 - d. The presence of blood in the airway device during its removal
2. The effectiveness of caruncular single peribulbar puncture technique in eye surgery after 10 min is approximately:
 - a. 90%
 - b. 80%
 - c. 70%
 - d. 60%
3. Phantom limb pain:
 - a. Is less frequent in elderly versus pediatric patients
 - b. Does not depend on the level of amputation
 - c. Is gender-dependent
 - d. Is neuropathic in origin
4. The following peripheral factors are related with the occurrence of phantom limb pain, except for:
 - a. Sympathetic activation
 - b. Neuroplasticity
 - c. Selective loss of C-fibers
 - d. Development of non-functional connections
5. The systemic local anesthetic toxicity following an inadvertent injection may be life threatening. Thus, a test dose is mandatory for preventing such event. The recommended drug is:
 - a. Lidocaine 2% with epinephrine
 - b. 100 mcg of fentanyl
 - c. 15 mg of 0.5% bupivacaine
 - d. 15 mg lidocaine 0.5% plus 25 mg epinephrine

[☆] Please cite this article as: Sanabria FR. Preguntas y respuestas. Rev Colomb Anestesiología. 2014;42:136–137.

* Correspondence to: Calle 119 # 7 – 75, Bogotá, Colombia.

E-mail address: raffanmago@gmail.com

6. Super morbid obesity is a patient with a BMI greater than:
 - a. 50 k/m²
 - b. 60 k/m²
 - c. 65 k/m²
 - d. 70 k/m²
7. Which of the following risk factors increase postoperative morbidity/mortality in the obese patient:
 - a. Female gender
 - b. Over 65 years old
 - c. BMI above 40
 - d. Asthma
8. In trauma and cardiovascular surgery, lactate and base deficit are both biomarkers that should be quantified very early on and in a serial mode, since they are both independent predictors of mortality within the first 48 h. Which of the following statements is false:
 - a. In the presence of mitochondrial oxygen, 36 ATPs are produced per glucose molecule during oxidative phosphorylation
 - b. Excess pyruvate converts into lactate via the lactic dehydrogenase enzyme that produces 2 ATP molecules
 - c. 2 categories of lactic acidosis are currently described: type A in the absence of tissue hypoxia and type B when lactic acidosis occurs with tissue hypoxia
 - d. Salicylate poisoning may result in lactic acidosis type B
9. All of the following are true with regard to lactate and base deficit measurements, except for:
 - a. The initial lactate value and its early clearance up to 9 h are independent early (less than 48 h) mortality factors
 - b. Lactate clearance enables the evaluation of the resuscitation process and quantification of the primary lesion
 - c. Regardless of the systolic pressure number, lactate levels of young patients admitted to the emergency room following blunt or penetrating trauma are predictive of the risk of receiving more than 6 red blood cell units in the first 24 h
 - d. Base deficit and lactate are correlated in hemorrhagic shock. The former is a sound indicator of the actual effective circulating volume deficit
10. Which of the following factors is not a predictor for difficult airway of the obese patient
 - a. Neck circumference (>40 cm)
 - b. Obstructive sleep apnea syndrome
 - c. Cervical mobility
 - d. Bite test

Reference

Raffan F. Preguntas y Respuestas. Rev Colomb Anestesiología. 2013;XLI(1);70-1.

Answers

- 1.b
- 2.a
- 3.d
- 4.b
- 5.b
- 6.b
- 7.d
- 8.c
- 9.c
- 10.a